arXiv:1201.3285v2 [math.RA] 12 Feb 2012

# Preserving meets in meet-dense poset completions

Robert Egrot

February 14, 2012

**Abstract**

Defining $P^*$ to be the complete lattice of up-sets (ordered by reverse inclusion) of a poset $P$ we give necessary and sufficient conditions on $\mathscr{S} \subseteq P^*$ for $P$ to admit a meet-completion $e\colon P \to Q$ where $e(\bigwedge S) = \bigwedge e[S]$ if and only if $S \in \mathscr{S}$. We show that, given $\mathscr{S}$ satisfying these conditions, the set $\mathbf{M}_{\mathscr{S}}$ of these completions is a topped weakly lower semimodular lattice when ordered by embedding lifting the identity on $P$. In particular, when $P$ is finite $\mathbf{M}_{\mathscr{S}}$ is a lower semimodular lattice, and a lower bounded homomorphic image of a free lattice. We provide an example of a countably infinite poset and a set $\mathscr{S} \subseteq P^*$ where $\mathbf{M}_{\mathscr{S}} \neq \emptyset$ has no bottom element.

## 1 Introduction

It is often desirable to embed a poset $P$ into a complete lattice $L$, preferably where the elements of $L$ are expressible in terms of the embedded copy of $P$. A famous example of this is Dedekind's construction of the reals from the rationals with cuts, a process generalized to posets by MacNeille [7], another being the generalization of the canonical extension construction (first defined for Boolean algebras [6]) to posets [1].

Here we are concerned with meet-completions, that is, completions where each element is expressible as an arbitrary meet of some subset of the original poset. It is well known (see proposition 3.1) that a meet-completion must preserve all existing joins from the original poset, and the fundamental observation here is that a meet-completion is characterized by the subsets of $P$ to which it assigns the same infimum (meet) (proposition 2.17). In particular, under certain circumstances (examined in section 3) we may choose a meet-completion of a poset that preserves specific existing meets and destroys all others.

Given a poset $P$, we define a condition we call *regularity* (definition 3.4) on a set $\mathscr{S} \subseteq P^*$ (where $P^*$ is the set of all up-sets of $P$) and show that it is equivalent to the property that every $S \in \mathscr{S}$ has an infimum in $P$ and there is at least one meet-completion of $P$ preserving all and only the meets of the sets in $\mathscr{S}$. In fact, we shall see that the set $\mathbf{M}_{\mathscr{S}}$ of these meet-completions forms a topped lattice (ordered via embedding lifting the identity on $P$) that is, in general, not bottomed (example 3.15). A necessary and sufficient condition on

$\mathscr{S}$ for $\mathbf{M}_{\mathscr{S}}$ to have a bottom element, and thus be a complete lattice, is given (proposition 3.13).

It has been shown [8] that every finite interval of the set of all join-completions of a poset $P$ (considered as a complete lattice ordered by embedding lifting the identity on $P$) is upper semimodular, an upper bounded homomorphic image of a free lattice, and meet-semidistributive. In section 4 we adapt and apply these results to show that $\mathbf{M}_{\mathscr{S}}$ is always weakly lower semimodular, and that when $P$ is finite $\mathbf{M}_{\mathscr{S}}$ is also lower semimodular, a lower bounded homomorphic image of a free lattice, and join-semidistributive.

Note that there is a well known duality between meet-completions of $P$ and certain closure operators on $P^*$, which we explain in section 2. For ease of exposition we phrase our results largely in terms of closure operators, implicitly appealing to this duality to justify our claims.

## 2 Preliminaries

Here we provide the basic definitions and notation we use in later sections, along with some background results.

**Definition 2.1** (embedding)**.** Given posets $P_1$ and $P_2$, a map $f\colon P_1 \to P_2$ is order preserving if $p \leq q \implies f(p) \leq f(q)$ for all $p, q \in P_1$, and it is an embedding if $p \leq q \iff f(p) \leq f(q)$ for all $p, q \in P_1$.

**Definition 2.2** (meet-dense)**.** Given a poset $Q$ and $P \subseteq Q$, we say $P$ is meet-dense in $Q$, or equivalently that $Q$ is a meet-dense extension of $P$, if whenever $q_1, q_2 \in Q$ with $q_1 \not\leq q_2$ there is $p \in P$ with $p \geq q_2$ and $p \not\geq q_1$.

**Definition 2.3** (meet-completion)**.** Given a poset $P$, a complete lattice $Q$, and an embedding $e\colon P \to Q$, we say $e\colon P \to Q$ is a meet-completion of $P$ if $e[P]$ is meet-dense in $Q$.

**Lemma 2.4.** *If $e_1\colon P \to Q_1$ and $e_2\colon P \to Q_2$ are meet-completions of $P$ and $g\colon Q_1 \to Q_2$ is an isomorphism such that $g \circ e_1 = e_2$, then $g$ is unique with this property.*

*Proof.* Suppose $h$ is another such isomorphism, then for all $p \in P$, and for all $q \in Q$, $e_1(p) \geq q \iff g \circ e_1(p) \geq g(q) \iff h \circ e_1(p) \geq h(q)$, and $g \circ e_1(p) \geq g(q) \iff e_2(p) \geq g(q)$, and similar for $h$, so $\{p \in P : e_2(p) \geq g(q)\} = \{p \in P : e_2(p) \geq h(q)\}$ and thus by meet-density we are done. ☐

**Definition 2.5** ($P^*$)**.** If $P$ is a poset define $P^*$ to be the complete lattice of up-sets (including $\emptyset$) of $P$ ordered by reverse inclusion.

**Definition 2.6** (closure operator)**.** Given a poset $P$ a closure operator on $P$ is a map $\Gamma\colon P \to P$ such that

1. $p \leq \Gamma(p)$ for all $p \in P$,
2. $p \leq q \implies \Gamma(p) \leq \Gamma(q)$ for all $p, q \in P$, and

3. $\Gamma(\Gamma(p)) = \Gamma(p)$ for all $p \in P$.

Following [2] we say a closure operator $\Gamma$ on $P^*$ or ${P^*}^\delta$ is *standard* when $\Gamma_e(p^\uparrow) = p^\uparrow$ for all $p \in P$ (here we use $\delta$ to signify the order dual).

**Lemma 2.7.** *$\iota\colon P \to P^*$ defined by $\iota(p) = p^\uparrow$ is a meet-completion of $P$.*

*Proof.* That $\iota\colon P \to P^*$ is a meet-completion is well known and straightforward to prove. Note though that $\iota$ will not map the top element of $P$ (if it exists) to the top element of $P^*$, as the top element of $P^*$ will be $\emptyset$. $\square$

It is well known that a meet-completion $e\colon P \to Q$ of $P$ defines a standard closure operator $\Gamma_e\colon\ P^{*\delta} \to P^{*\delta}$ by $\Gamma_e(S) = \{p \in P : e(p) \geq \bigwedge e[S]\}$ (we take the dual of $P^*$ as otherwise condition 1 of definition 2.6 fails). In this case $Q$ is isomorphic to the lattice $\Gamma_e[P^*]$ of $\Gamma_e$-closed subsets of $P^*$ (note we are purposefully taking $P^*$ here rather than ${P^*}^\delta$ as we want to order by reverse inclusion, this is technically an abuse of notation as $\Gamma_e$ is originally defined on ${P^*}^\delta$, but as these structures have the same carrier hopefully our meaning is clear) via the map $h_e\colon Q \to \Gamma_e[P^*]$ defined by $h_e(q) = \{p \in P : e(p) \geq q\}$. Conversely, whenever $\Gamma$ is a standard closure operation on $P^{*\delta}$, $\Gamma$ induces a meet-completion $e_\Gamma\colon P \to \Gamma[P^*]$ defined by $e_\Gamma(p) = p^\uparrow$. For $S \in P^*$ we have $\Gamma_{e_\Gamma}(S) = \{p : p^\uparrow \subseteq \bigwedge S\} = \{p : p^\uparrow \subseteq \Gamma(S)\} = \Gamma(S)$, so $\Gamma_{e_\Gamma} = \Gamma$, and, for all $p \in P$, $e_{\Gamma_e}(p) = p^\uparrow = h_e \circ e(p)$ so the following commutes:

$$\begin{tikzcd}
P \arrow[r, "e"] \arrow[d, "e_{\Gamma_e}"'] & Q \arrow[dl, "h_e"] \\
\Gamma_e[P^*] &
\end{tikzcd}$$

We state the results of the preceeding discussion as a theorem.

**Theorem 2.8.** *If $e\colon P \to Q$ is a meet-completion then there is a unique isomorphism such that the following commutes:*

$$\begin{tikzcd}
P \arrow[r, "e"] \arrow[d, "e_{\Gamma_e}"'] & Q \\
\Gamma_e[P^*] \arrow[ur, leftrightarrow, "\cong"'] &
\end{tikzcd}$$

*Moreover, if $e_1\colon P \to Q_q$ and $e_2\colon P \to Q_2$ are meet-completions such that there is an isomorphism $h\colon Q_1 \to Q_2$ with $h \circ e_1 = e_2$ then $\Gamma_{e_1} = \Gamma_{e_2}$.*

*Proof.* The existence of the required isomorphisms has been established, and uniqueness follows from lemma 2.4. If $h\colon Q_1 \to Q_2$ with $h \circ e_1 = e_2$ then $\Gamma_{e_2} = \{p \in P : e_2(p) \geq \bigwedge e_2[S]\} = \{p \in P : h \circ e_1(p) \geq \bigwedge h \circ e_1[S]\} = \{p \in P : h \circ e_1(p) \geq h(\bigwedge e_1[S])\} = \{p \in P : e_1(p) \geq \bigwedge e_1[S]\} = \Gamma_{e_1}(S)$. $\square$

**Definition 2.9** (lift)**.** Given posets $P_1, P_2, Q_1$, and $Q_2$, embeddings $e_1\colon P_1 \to Q_1$ and $e_2\colon P_2 \to Q_2$, and order preserving $f\colon P_1 \to P_2$, we say the lift of $f$ along $e_1$ and $e_2$ (or just the lift of $f$ when $e_1$ and $e_2$ are clear from context) is an order preserving map $f'\colon Q_1 \to Q_2$ such that the following commutes:

$$\begin{tikzcd} P_1 \arrow[r, "e_1"] \arrow[d, "f"'] & Q_1 \arrow[d, "f'"] \\ P_1 \arrow[r, "e_2"'] & Q_2 \end{tikzcd}$$

The meet-completions of a poset $P$ can be characterized up to isomorphism by a generalized concept of meet-primality, which we explain below. In particular the algebra $P^*$ is the unique (up to isomorphism lifting the identity on $P$) meet-completion $P$ whose set of completely meet-prime elements is the embedded image of $P$ (corollary 2.16).

**Definition 2.10** (relatively meet-prime)**.** Let $L$ be a lattice, let $\mathscr{X} \subseteq \wp(L)$, and let $a \in L$. Then $a$ is meet-prime relative to $\mathscr{X}$ if whenever $X \in \mathscr{X}$ and $a \geq \bigwedge X$ there is $x \in X$ with $a \geq x$.

**Definition 2.11** ($\mathscr{X}_\Gamma$)**.** Given a closure operator $\Gamma\colon P^{*\delta} \to P^{*\delta}$ define $\mathscr{X}_\Gamma = \{X \in \wp(\Gamma[P^*]) : \Gamma(\bigcup X) = \bigcup X\}$.

**Lemma 2.12.** *Let $e\colon P \to Q$ be a meet-completion, then the set of elements of $Q$ that are meet-prime relative to $\mathscr{X} = \{X \in \wp(Q) : h_e[X] \in \mathscr{X}_{\Gamma_e}\}$ is precisely $e[P]$.*

*Proof.* It is given that $Q$ is a complete lattice into which $P$ embeds meet-densely via $e$, so we show that the set of elements of $Q$ that are meet-prime relative to $\mathscr{X}_{\Gamma_e}$ is precisely $e[P]$. Let $q \in Q$, then $q$ is meet-prime relative to $\mathscr{X}$ if and only if, for all $X \in \mathscr{X}$, $q \geq \bigwedge X \implies q \geq x$ for some $x \in X$, which is equivalent to saying that $h_e(q) \subseteq \bigcup h_e[X] \implies h_e(q) \subseteq h_e(x)$ for some $x \in X$. If $q \in e[P]$ then this clearly holds as $h_e(e(p)) = p^\uparrow$ for all $p \in P$. Conversely, suppose $q \notin e[P]$ and let $X = \{e(p) : p \in h_e(q)\}$, then $\Gamma_e(\bigcup X) = \bigcup X$, and clearly $h_e(q) = \bigcup h_e[X]$, but $h_e(q) \not\subseteq h_e(x)$ for all $x \in X$, and thus $q$ cannot be meet-prime relative to $\mathscr{X}$. ☐

**Lemma 2.13.** *If $e_1\colon P \to Q_1$ and $e_2\colon P \to Q_2$ are meet-completions of $P$ then $\mathscr{X}_{\Gamma_{e_1}} = \mathscr{X}_{\Gamma_{e_2}} \iff \Gamma_{e_1} = \Gamma_{e_2}$.*

*Proof.* Suppose $C \in \Gamma_{e_1}[P^*] \setminus \Gamma_{e_2}[P^*]$, then $\Gamma_{e_1}(\bigcup\{C\}) = \Gamma_{e_1}(C) = C = \bigcup\{C\}$, so $C \in \mathscr{X}_{\Gamma_{e_1}}$, but $\Gamma_{e_2}(C) \supset C$ so $\{C\} \notin \mathscr{X}_{\Gamma_{e_2}}$. The converse is automatic. ☐

**Lemma 2.14.** *If $e_1\colon P \to Q_1$ and $e_2\colon P \to Q_2$ are meet-completions of $P$ then $Q_1 \cong Q_2$ via an isomorphism lifting the identity map on $P$ if and only if $\Gamma_{e_1} = \Gamma_{e_2}$.*

*Proof.* If $\Gamma_{e_1} = \Gamma_{e_2}$ then the required isomporphism exists by theorem 2.8. Conversely, the existence of such an isomorphism demands that $\Gamma_{e_1} = \Gamma_{e_2}$, by definition of $\Gamma_e$. $\square$

**Lemma 2.15.** *If $e\colon P \to Q$ is a meet-completion such that the set of completely meet-prime elements of $Q$ is precisely $e[P]$, then $\mathscr{X}_{\Gamma_e} = \wp(P^*)$.*

*Proof.* Suppose the set of completely meet-prime elements of $Q$ is precisely $e[P]$ and let $X \in \wp(P^*)$. Define $X' = \{p^\uparrow : p \in \bigcup X\}$, then $\bigcup X' = \bigcup X$ and $\Gamma_e(\bigcup X') = \Gamma_e(\bigcup X)$. Now, $p \in \Gamma_e(\bigcup X') \iff e(p) \geq \bigwedge\{e(p') : p'^\uparrow \in X'\}$, so $p \in \Gamma_e(\bigcup X') \implies e(p) \geq e(p')$ for some $p' \in \bigcup X \implies p \in \bigcup X$, and thus $\Gamma_e(\bigcup X) = \bigcup X$. It follows that $\Gamma_e(S) = S$ for all $S \in P^*$, and $\mathscr{X}_{\Gamma_e} = \wp(P^*)$. $\square$

**Corollary 2.16.** *$P^*$ can be characterized abstractly as the unique (up to isomorphism lifting the identity on $P$) meet-completion of $P$ whose set of completely meet-prime elements is the embedded image of $P$.*

*Proof.* $P^*$ corresponds to the closure operator on $P^{*\delta}$ that is the identity map, and for this closure operator the set $\mathscr{X}_\Gamma$ is $\wp(P^*)$. That the completely meet-prime elements of $P^*$ is $\iota[P]$ follows from lemma 2.12. If $e\colon P \to Q$ is a meet-completion of $P$ whose set of completely meet-prime elements is $e[P]$, then$\mathscr{X}_{\Gamma_e} = \wp(P^*)$ by lemma 2.15 , and so lemmas 2.13 and 2.14 together say required isomorphism exists. $\square$

We make the simple observation that a meet-completion is defined (up to isomorphism lifting the identity on $P$) by the subsets of $P$ to which it assigns the same infimum:

**Proposition 2.17.** *If $e_1\colon P \to Q_1$ and $e_2\colon P \to Q_2$ are meet-completions then there is a unique isomorphism $h\colon Q_1 \to Q_2$ lifting the identity on $P$ if and only if for all $S, T \subseteq P$ we have $\bigwedge e_1[S] = \bigwedge e_1[T] \iff \bigwedge e_2[S] = \bigwedge e_2[T]$ .*

*Proof.* That the existence of a unique isomorphism $h\colon Q_1 \to Q_2$ with $h \circ e_1 = e_2$ implies for all $S, T \subseteq P$ we have $\bigwedge e_1[S] = \bigwedge e_1[T] \iff \bigwedge e_2[S] = \bigwedge e_2[T]$ is trivial. For the converse note that to say $\bigwedge e_1[S] = \bigwedge e_1[T] \iff \bigwedge e_2[S] = \bigwedge e_2[T]$ is equivalent to saying that $\Gamma_{e_1}(S) = \Gamma_{e_1}(T) \iff \Gamma_{e_2}(S) = \Gamma_{e_2}(T)$. Let $X \notin \mathscr{X}_{\Gamma_{e_1}}$, then $\Gamma_{e_1}(\bigcup X) = C = \Gamma_{e_1}(C) \supset \bigcup X$ for some $C \in P^*$, so $\Gamma_{e_2}(\bigcup X) = \Gamma_{e_2}(C) \supset \bigcup X$, and thus $X \notin \mathscr{X}_{\Gamma_{e_2}}$ and so we must have $\mathscr{X}_{\Gamma_{e_2}} \subseteq \mathscr{X}_{\Gamma_{e_1}}$. By symmetry we also have $\mathscr{X}_{\Gamma_{e_1}} \subseteq \mathscr{X}_{\Gamma_{e_2}}$ and the result follows from lemmas 2.13 and 2.14. $\square$

We can say more about relationship between meet completions and standard closure operators:

**Definition 2.18** ($\mathbb{S}_P$)**.** Define $\mathbb{S}_P$ to be the complete lattice of standard closure operators on $P^{*\delta}$ (ordered pointwise, i.e. $\Gamma_1 \leq \Gamma_2 \iff \Gamma_1(S) \leq \Gamma_2(S) \iff \Gamma_1(S) \subseteq \Gamma_2(S)$ for all $S \in P^*$).

**Definition 2.19** ($\mathbf{M}_P$)**.** Define $\mathbf{M}_P$ to be the set of meet completions of $P$, ordered by defining $e_1 \colon P \to Q_1 \leq e_2 \colon P \to Q_2 \iff$ there is an embedding $e \colon Q_1 \to Q_2$ lifting the identity on $P$. Since we have $e_1 \leq e_2$ and $e_2 \leq e_1 \iff Q_1 \cong Q_2$ we can consider $\mathbf{M}_P$ to be a poset.

**Lemma 2.20.** *For all* $\Gamma_1, \Gamma_2 \in \mathbb{S}_P$*,* $\Gamma_1 \leq \Gamma_2 \iff \Gamma_2[P^*]$ *embeds into* $\Gamma_1[P^*]$ *via a map lifting the identity on* $P$*.*

*Proof.* Suppose $\Gamma_1 \leq \Gamma_2$ and define $f \colon \Gamma_2[P^*] \to \Gamma_1[P^*]$ by $f(C) = \Gamma_1(C) = C$. Then $f$ is well defined as $\Gamma_1 \leq \Gamma_2$, and certainly is an embedding lifting the identity on $P$. Conversely, for all $S \in P^*$ and for all $p \in P$, $p \in e(\Gamma_2(S)) \iff p^\uparrow \geq e(\Gamma_2(S)) \iff p^\uparrow \geq \Gamma_2(S) \iff p \in \Gamma_2(S)$, so $e(\Gamma_2(S)) = \Gamma_2(S)$, and thus $\Gamma_2(S)$ is $\Gamma_1$-closed. Since $\Gamma_1(S)$ must be the smallest $\Gamma_1$-closed set containing $S$ we must have $\Gamma_1(S) \subseteq \Gamma_2(S)$. $\square$

**Theorem 2.21.** $\mathbf{M}_P$ *and* $\mathbb{S}_P$ *are dually isomorphic.*

*Proof.* This follows easily from theorem 2.8 and lemma 2.20. $\square$

Finally we give an example of two meet-completions of a poset $P$ that are isomorphic but with no isomorphism lifting the identity on $P$ (example 2.22).

**Example 2.22.** Let $P = \{a, b, c\}$ be the three element antichain, and let $Q_1$ and $Q_2$ be as in figure 1. Suppose the image of $P$ in $Q_1$ and $Q_2$ is as marked, then $Q_1$ and $Q_2$ are clearly order isomorphic, but there is no such isomorphism that is a lift of the identity on $P$.



Figure 1: Meet completions of $P$ that are isomorphic but not via any isomorphism lifting the identity on $P$

# 3 The preservation of meets

It is well known that meet-completions preserve all existing joins (see proposition 3.1), existing meets however are not necessarily preserved, though subject to certain constraints we can construct meet-completions preserving only specific meets, in a sense we shall make precise in this section.

**Proposition 3.1.** *Let* $e \colon P \to Q$ *be a meet-completion of* $P$*, then for all* $S, T \subseteq P$*,* $\bigvee S = \bigvee T \implies \bigvee e[S] = \bigvee e[T]$*. Conversely, if either* $\bigvee S$ *or* $\bigvee T$ *exist in* $P$ *then* $\bigvee e[S] = \bigvee e[T] \implies$ *they both exist and are equal.*

*Proof.* Suppose $\bigvee e[S] \not\leq \bigvee e[T]$, then by meet-density there is $p \in P$ with $e(p) \geq \bigvee e[T]$ and $e(p) \not\geq \bigvee e[S]$. So $p \geq t$ for all $t \in T$, and there is $s \in S$ with $p \not\geq s$, and so $p \geq \bigvee T$ but $p \not\geq \bigvee S$ and so $\bigvee S \not\leq \bigvee T$. The result follows by symmetry. Conversely, let $\bigvee S = p \in P$ and suppose $\bigvee e[S] = \bigvee e[T]$. Then $e(p) = \bigvee e[S] = \bigvee e[T]$, so $e(p) \leq e(t)$ for all $t \in T$, and as $e$ is an embedding this means $p \leq t$ for all $t \in T$. If $z \leq t$ for all $t \in T$ then $e(z) \leq e(t)$ for all $t \in T$, but then $e(z) \leq \bigvee e[t] = \bigvee e[S] = e(p)$, and so $z \leq p$, and thus $\bigvee T = p = \bigvee S$ as required. □

**Lemma 3.2.** *Let* $e\colon P \to Q$ *be an embedding, let* $S \in P^*$*, and suppose* $\bigwedge S = p$ *in* $P$*. Then* $\bigwedge e[S] = e(p)$ *in* $Q$ *if and only if* $\Gamma_e(S) = p^\uparrow$*.*

*Proof.* Since $e(p)$ is clearly a lower bound for $e[S]$, $\bigwedge e[S] = e(p)$ if and only if whenever $q$ is a lower bound for $e[S]$ we have $q \leq e(p)$, but this occurs if and only if whenever $C \in \Gamma_e[P^*]$ and $C \leq \Gamma_e(s)$ for all $s \in S$ we have $C \leq \Gamma_e(p)$, but this is just saying that whenever $C \in \Gamma_e[P^*]$ and $S \subseteq C$ we have $p \in C$, which in turn is equivalent to saying that $p \in \Gamma_e(S)$. Since $\Gamma_e(p^\uparrow) = p^\uparrow$ and $S \subseteq p^\uparrow$ we have $p \in \Gamma_e(S) \iff \Gamma_e(S) = p^\uparrow$. □

**Definition 3.3** ($\mathscr{S}$-regular)**.** Given $\mathscr{S} \subseteq P^*$, we say a meet-completion $e\colon P \to Q$ is $\mathscr{S}$-regular if $\bigwedge S$ is defined in $P$ and $e(\bigwedge S) = \bigwedge e[S]$ for all $S \in \mathscr{S}$. If $\mathscr{S}$ contains every set $S$ for which $\bigwedge S$ is defined in $P$, then we say that an $\mathscr{S}$-regular meet-completion is *regular*.

**Definition 3.4** (regular)**.** Let $P$ be a poset, then $\mathscr{S} \subseteq P^*$ is regular if

1. $p^\uparrow \in \mathscr{S}$ for all $p \in P$,
2. $\bigwedge S$ exists in $P$ for all $S \in \mathscr{S}$, and
3. whenever $T \in P^* \setminus \mathscr{S}$, there is $T' \in P^*$ with $T \subseteq T'$, $p < T \implies p < T'$ for all $p \in P$, and for all $S \in \mathscr{S}$, $S \subseteq T' \implies \bigwedge S \in T'$.

This definition appears rather technical, but the intuition behind it is quite simple. If $e\colon P \to Q$ is a meet-completion then, for every $q \in Q$, $\{p \in P : e(p) \geq q\}$ is clearly an up-set, and thus every meet-completion can be thought of as being based on a subset of $P^*$ (this is implicit in our definition via closure operators). A connection between preservation of finite meets and the closure of the up-sets it is based on under certain finite infima is known (see for example [4, Proposition 6.10]), and it is not hard to show that this extends to arbitrary meets too. The key point is that a meet-completion will preserve the infimum of a set $S$ if and only if every up-set on which the completion is based contains that infimum whenever it contains $S$. Condition 3 of definition 3.4 essentially demands that every set $T$ whose infimum we wish to destroy can be contained in an up-set which also does not contain the infimum of $T$ but does not interfere with the preservation of the infima we *do* wish to preserve.

**Definition 3.5** ($\mathscr{S}$-closure)**.** If $\mathscr{S} \subseteq P^*$ we say a standard closure operator $\Gamma\colon {P^*}^\delta \to {P^*}^\delta$ is an $\mathscr{S}$-closure if for all $S \in P^*$, $\Gamma(S) = p^\uparrow$ for some $p \in P \iff S \in \mathscr{S}$.

**Definition 3.6** ($\mathcal{F}_{\mathscr{S}}$, $\Gamma_{\mathscr{S}}$)**.** If $\mathscr{S} \subseteq P^*$ is regular define $\mathcal{F}_{\mathscr{S}} = \{f \in P^* : S \subseteq f \implies \bigwedge S \in f$ for all $S \in \mathscr{S}\}$, and define $\Gamma_{\mathscr{S}} \colon P^{*^\delta} \to P^{*^\delta}$ by $\Gamma_{\mathscr{S}}(S) = \bigcap\{f \in \mathcal{F}_{\mathscr{S}} : S \subseteq f\}$.

**Lemma 3.7.** *If $\mathscr{S} \subseteq P^*$ is regular then $\Gamma_{\mathscr{S}}$ is an $\mathscr{S}$-closure.*

*Proof.* As $\mathcal{F}_{\mathscr{S}}$ is closed under arbitrary intersections it's easy to see that $\Gamma_{\mathscr{S}}$ is a standard closure operator on $P^{*^\delta}$, and if $S \in \mathscr{S}$ then $\Gamma_{\mathscr{S}}(S) = p^\uparrow$ for some $p \in P$ by definition of $\mathcal{F}$. If $T \in P^* \setminus \mathscr{S}$ then by definition of regularity there is $T' \in \mathcal{F}_{\mathscr{S}}$ with $T \subseteq T' \subset p^\uparrow$ whenever $p$ is a lower bound for $T$, and since we must have $\Gamma_{\mathscr{S}}(T) \subseteq \Gamma_{\mathscr{S}}(T') = T' \subset p^\uparrow$ for all lower bounds $p$ of $T$ we cannot have $\Gamma_{\mathscr{S}}(T) = p^\uparrow$ for any $p \in P$, as $\Gamma_{\mathscr{S}}(T) = p^\uparrow \implies p < T$. ☐

**Proposition 3.8.** *Let $P$ be a poset, and let $\mathscr{S} \subseteq P^*$. Then $\mathscr{S}$ is regular if and only if there exists an $\mathscr{S}$-closure.*

*Proof.* Suppose first that $\Gamma$ is an $\mathscr{S}$-closure, then by definition we must have $p^\uparrow \in \mathscr{S}$ for all $p \in P$, and if $S \in \mathscr{S}$ then $\Gamma(S) = p^\uparrow$ for some $p \in P$. If $p$ is not the greatest lower bound of $S$ there must be $p' \in P$ with $p' \leq S$ and $p' \not\leq p$. But then $S \subseteq \Gamma(p'^\uparrow)$ and thus $S \subseteq \Gamma(p'^\uparrow) \cap \Gamma(p^\uparrow) \subset p^\uparrow$, which contradicts the facts that the intersection of $\Gamma$-closed sets is closed and $\Gamma(S)$ must be the smallest $\Gamma$-closed set containing $S$. Now suppose $T \in P^* \setminus \mathscr{S}$ and let $p \leq T$, then by definition of $\mathscr{S}$-closure $\Gamma(T) \neq p^\uparrow$, so $T \subseteq \Gamma(T) \subset p^\uparrow$, and for all $S \in \mathscr{S}$ we have $S \subseteq \Gamma(T) \implies \Gamma(S) \subseteq \Gamma(T) \implies \bigwedge S \in \Gamma(T)$ as required. The converse follows trivially from lemma 3.7. ☐

**Corollary 3.9.** *Let $P$ be a poset and let $\mathscr{S} \subseteq P^*$, then $\mathscr{S}$ is regular if and only if there is at least one $\mathscr{S}$-regular meet-completion.*

*Proof.* If $\mathscr{S}$ is regular just take $\Gamma_{\mathscr{S}}$ and lemma 3.2 gives the result. Conversely, an $\mathscr{S}$-regular meet-completion $e \colon P \to Q$ gives rise to a standard closure operator $\Gamma_e$, which lemma 3.2 says must be an $\mathscr{S}$-closure. The result then follows from proposition 3.8. ☐

**Lemma 3.10.** *Let $\mathscr{S} \subseteq P^*$ be regular, and let $\Gamma$ be an $\mathscr{S}$-closure. Then, for all $T \in P^{*^\delta}$, $\Gamma_{\mathscr{S}}(T) \leq \Gamma(T)$.*

*Proof.* Let $T \in P^{*^\delta}$, and let $S \in \mathscr{S}$, then $S \subseteq \Gamma(T) \implies \Gamma(S) \subseteq \Gamma(T) \iff \bigwedge S \in \Gamma(T)$, so $\Gamma(T) \in \mathcal{F}_{\mathscr{S}}$ and thus $\Gamma_{\mathscr{S}}(T) \subseteq \Gamma(T)$ and we have the result. ☐

**Lemma 3.11.** *If $\Gamma_1$ and $\Gamma_2$ are $\mathscr{S}$-closures then $\Gamma_2 \circ \Gamma_1$ is also an $\mathscr{S}$-closure.*

*Proof.* For all $S \in P^*$, $\Gamma_1(\Gamma_2(S)) = p^\uparrow \iff \Gamma_2(S) \in \mathscr{S} \iff \Gamma_2(\Gamma_2(S)) = p^\uparrow \iff \Gamma_2(S) = p^\uparrow$ for some $p \in P \iff S \in \mathscr{S}$. ☐

**Theorem 3.12.** *If $P$ is a poset and $\mathscr{S} \subseteq P^*$ is regular then the set $\mathbb{S}_{\mathscr{S}}$ of $\mathscr{S}$-closures is a lattice with bottom element $\Gamma_{\mathscr{S}}$ when ordered pointwise (i.e. $\Gamma_1 \leq \Gamma_2 \iff \Gamma_1(S) \leq \Gamma_2(S) \iff \Gamma_1(S) \subseteq \Gamma_2(S)$ for all $S \in P^*$). Moreover, arbitrary non-empty meets are defined in $\mathbb{S}_{\mathscr{S}}$.*

*Proof.* Lemma 3.10 tells us $\Gamma_{\mathscr{S}}$ is mimimal in the set of $\mathscr{S}$-closures, and the meet operation on $\mathbb{S}_{\mathscr{S}}$ is defined by set intersection. For $\Gamma_1, \Gamma_2 \in \mathbb{S}_{\mathscr{S}}$, $\Gamma = \Gamma_2 \circ \Gamma_1$ is an upper bound for $\{\Gamma_1, \Gamma_2\}$ in $\mathbb{S}_{\mathscr{S}}$ (using lemma 3.11), so by closure under arbitrary meets the join of $\Gamma_1$ and $\Gamma_2$ must be defined in $\mathbb{S}_{\mathscr{S}}$. ☐

In light of theorem 3.12, $\mathbb{S}_{\mathscr{S}}$ will be a complete lattice if and only if it has a top element, which it may not (see example 3.15). Proposition 3.13 gives a necessary and sufficient condition for $\mathbb{S}_{\mathscr{S}}$ to have a top, and corollary 3.14 gives a simpler sufficient condition.

**Proposition 3.13.** *Let $P$ be a poset and let $\mathscr{S} \subseteq P^*$ be regular. For each $T \in P^* \setminus \mathscr{S}$ define $\bar{T} = \{T' \in \mathcal{F}_{\mathscr{S}} : T \subseteq T'$, and $p < T \implies p < T'$ for all $p \in P\}$. Then $\mathbb{S}_{\mathscr{S}}$ has a top element if and only if for each $T \in P^* \setminus \mathscr{S}$ there is $T^m \in \bar{T}$ such that for every $Y \subseteq \bigcup\{\bar{U} : U \in P^* \setminus \mathscr{S}\}$ with $Y \cap \bar{U} \neq \emptyset$ for all $U \in P^* \setminus \mathscr{S}$ we have $\bigcap\{f \in Y : T \subseteq f\} \subseteq T^m$.*

*Proof.* Suppose such $T^m$ exist, and suppose wlog that for each $T$, $T^m$ is the smallest set satisfying the condition (this is possible as $\bar{T}$ is closed under intersection, and so is the subset satisfying our condition). Define $\mathcal{F}' = \{p^\uparrow : p \in P\} \cup \bigcup\{T^m : T \in P^* \setminus \mathscr{S}\}$, define $\mathcal{F}$ to be the closure of $\mathcal{F}'$ under arbitrary intersections, and define $\Gamma\colon {P^*}^\delta \to {P^*}^\delta$ by $\Gamma(U) = \bigcap\{f \in \mathcal{F} : U \subseteq f\}$. Then $\Gamma$ is clearly a closure operator on ${P^*}^\delta$ and $\Gamma(p^\uparrow) = p^\uparrow$ for all $p \in P$. If $T \notin \mathscr{S}$ then $\Gamma(T) = T^m$ (by minimality of $T^m$), and $T^m \neq p^\uparrow$ for all $p \in P$. If $S \in \mathscr{S}$ then, for all $f \in \mathcal{F}'$, $S \subseteq f \implies (\bigwedge S)^\uparrow \subseteq f$, so $\Gamma(S) = (\bigwedge S)^\uparrow$, and thus $\Gamma$ is an $\mathscr{S}$-closure.

If $\Gamma'$ is another $\mathscr{S}$-closure then $\Gamma(S) = (\bigwedge S)^\uparrow = \Gamma'(S)$ for all $S \in \mathscr{S}$. Define $Y = \{\Gamma'(U) : U \in P^* \setminus \mathscr{S}\}$, then, since from the proof of proposition 3.8 we know that $\Gamma'(U) \in \bar{U}$ for all $U \in P^* \setminus \mathscr{S}$, we have $Y \subseteq \bigcup\{\bar{U} : U \in P^* \setminus \mathscr{S}\}$ and $Y \cap \bar{U} \neq \emptyset$ for all $U \in P^* \setminus \mathscr{S}$. So, given $T \in P^* \setminus \mathscr{S}$, we have $\Gamma'(T) = \bigcap\{f \in Y : T \subseteq f\} \subseteq T^m = \Gamma(T)$, and thus $\Gamma' \leq \Gamma$ as required.

Conversely, suppose $\Gamma$ is a top element for $\mathbb{S}_{\mathscr{S}}$, and let $Y$ be any subset of $\bigcup\{\bar{T} : T \in P^* \setminus \mathscr{S}\}$ such that $\bar{T} \cap Y \neq \emptyset$ for all $T \in P^* \setminus \mathscr{S}$. Then the closure under intersections of $\mathcal{F} = \{p^\uparrow : p \in P\} \cup Y$ defines an $\mathscr{S}$-closure, $\Gamma_Y$, and $\Gamma_Y(T) = \bigcap\{f \in Y : T \subseteq f\}$ for all $T \in P^* \setminus \mathscr{S}$. By maximality of $\Gamma$ we have $\Gamma_Y(T) \subseteq \Gamma(T)$ for all $T \in P^* \setminus \mathscr{S}$, which means $\bigcap\{f \in Y : T \subseteq f\} \subseteq \Gamma(T)$, so we can take $T^m = \Gamma(T)$. Moreover, if we define $Y' = \{\Gamma(T) : T \in P^* \setminus \mathscr{S}\}$ then $Y' \subseteq \bigcup\{\bar{U} : U \in P^* \setminus \mathscr{S}\}$ and $Y \cap \bar{U} \neq \emptyset$. Since $\Gamma(T) = \bigcap\{f \in Y' : T \subseteq f\}$ $T^m$ is also the smallest set satisfying the required condition. ☐

**Corollary 3.14.** *If $\bar{T}$ has at least one maximal element (ordering by inclusion) for all $T \in P^* \setminus \mathscr{S}$ then $\mathbb{S}_{\mathscr{S}}$ has a top element and thus is complete.*

*Proof.* Given $T \in P^* \setminus \mathscr{S}$ define $M_T \subseteq \bar{T}$ to be the set of maximal elements of $\bar{T}$, and define $T^m = \bigcap M_T$. Then certainly $T^m \in \bar{T}$, and if we choose any $Y \subseteq \bigcup\{\bar{T} : T \in P^* \setminus \mathscr{S}\}$ with $Y \cap \bar{U} \neq \emptyset$ for all $U \in P^* \setminus \mathscr{S}$ we must have $m \in Y$ for all $m \in M_T$, as $Y \cap \bar{m} \neq \emptyset$ by choice of $Y$, and $\bar{m} = \{m\}$ by maximality of $m$. So $\bigcap\{f \in Y : T \subseteq f\} \subseteq \bigcap M_T = T^m$, and by proposition 3.13 we have the result. ☐

**Example 3.15.** Let $P' = \{p_n : n \in \omega\}$ be an antichain, let $P = P' \cup \{0\}$, where 0 is a a bottom element, and let $\mathscr{S} = \{p^\uparrow : p \in P\} \cup \{S \in P^* : |S| = |\omega|\}$ (it's easy to check that $\mathscr{S}$ is regular). Let $U = \{p_1, p_2\}$ and let $U' \in \bar{U}$. Since $U'$ must be finite we can choose $p \in P' \setminus U'$ and define $Y = \{f \in P^* : f$ is finite and $U' \cup \{p\} \subseteq f\}$. Then $Y \subseteq \bigcup\{\bar{T} : T \in P^* \setminus \mathscr{S}\}$, and $Y \cap \bar{T} \neq \emptyset$ for all $T \in P^* \setminus \mathscr{S}$, however, $\bigcap\{f \in Y : U \subseteq f\} \supseteq U' \cup \{p\} \supset U'$, so by proposition 3.13 $\mathbb{S}_{\mathscr{S}}$ has no top element.

## 4 The structure of $\mathbb{S}_{\mathscr{S}}$

We can say a little more about the structure of $\mathbb{S}_{\mathscr{S}}$, though first we require some preliminary definitions and results.

**Definition 4.1** (weakly lower/upper semimodular)**.** A poset $P$ is weakly lower semimodular if whenever $a, b, c \in P$ with $a \neq b$, $a \prec c$ and $b \prec c$ there is $d$ with $d \prec a$ and $d \prec b$ (where $\prec$ is the 'covers' relation defined by $x \prec y \iff x < y$ and there is no $z$ with $x < z < y$). Weakly upper semimodularity is defined dually

This definition appears as a definition for lower/upper semimodularity in [5]. In the special case where $P$ is a lattice these conditions are implied by lower/upper semimodularity (see definition 4.2 below) respectively, and in a dually strongly atomic lattice the converse also holds (a lattice is *strongly atomic* if whenever $a < b$ there is $p$ with $a \prec p \leq b$). Note that a finite lattice is always dually strongly atomic.

**Definition 4.2** (lower/upper semimodular)**.** A lattice $L$ is lower semimodular if for all $a, b, c \in L$, if $a \parallel b$ and $a < b < a \vee c$ then there is some $d \in L$ with $c \leq d < a \vee c$ and $a \vee (b \wedge d) = b$. Upper semimodularity is defined dually.

**Definition 4.3** (upper/lower bounded homomorphism)**.** A lattice homomorphism $h \colon K \to L$ is upper bounded if $\{b \in K : h(b) \leq a\}$ is either empty or has a greatest element for all $a \in L$. Lower bounded homomorphisms are defined dually.

Given a complete lattice $L$ we define $\mathbb{CL}(L)$ to be the complete lattice of closure operators on $L$ (ordered pointwise), and we define $\mathbb{SUB}_\wedge(L)$ to be the complete lattice of meet subsemilattices of $L$ (regarded as a meet semilattice) containing 1.

**Theorem 4.4** (originating in [9], reproduced here as it appears in [8])**.** *If $L$ is a complete lattice then $\mathbb{CL}(L)$ is dually isomorphic to $\mathbb{SUB}_\wedge(L)$.*

**Lemma 4.5** (this is lemma 2 of [8])**.** *Let $L$ be a complete lattice. For all $U, V \in \mathbb{SUB}_\wedge(L)$, if $U \prec V$ then $U = V \setminus \{a\}$ for some $a \in V$.*

**Corollary 4.6.** *If $S \subseteq \mathbb{SUB}_\wedge(L)$ has the following properties*

1. *the interval $[y, x] \subseteq S$ for all $y \leq x \in S$, and*

*2. $S$ is closed under finite meets (inherited from $\mathbb{SUB}_\wedge(L)$)*

*then $S$ is weakly lower semimodular.*

*Proof.* Let $U, V, W \in S$, and suppose $U \prec W$ and $V \prec W$. Then by lemma 4.5 $U = W\backslash\{a\}$ and $V = W\backslash\{b\}$ for some $a \neq b \in W$, so $U \wedge V = U \cap V = W\backslash\{a, b\}$, and so clearly $U \wedge V \prec U$ and $U \wedge V \prec V$ so we are done. ☐

**Proposition 4.7.** *If $\mathscr{S} \subseteq P^*$ is regular then $\mathbb{S}_\mathscr{S}$ is weakly upper semimodular.*

*Proof.* This follows easily from theorem 4.4, corollary 4.6, and the fact that $\mathbb{S}_\mathscr{S}$ is a lattice. ☐

When $P$ is finite we can go further, as in this case $\mathbb{S}_\mathscr{S}$ is a finite interval of $\mathbb{CL}(P^{*^\delta})$, and we can apply [8, theorem 6] (reproduced for convenience as theorem 4.9 below) directly to obtain the following:

**Theorem 4.8.** *If $P$ is finite, and if $\mathscr{S} \subseteq P^*$ is regular, then $\mathbb{S}_\mathscr{S}$ is upper semimodular, an upper bounded homomorphic image of a free lattice, and thus meet semidistributive.*

See e.g. [3, chapter 2] for further discussion of these ideas. The duality between $\mathbf{M}_P$ and $\mathbb{S}_P$ from theorem 2.21 restricted to $\mathbb{S}_\mathscr{S}$ gives dual results to those in this and the last section for $\mathbf{M}_\mathscr{S}$, the set of meet completions of $P$ preserving those, and only those, meets that are defined in $\mathscr{S}$.

**Theorem 4.9.** *[8, theorem 6] Let $L$ be a complete lattice. Then every finite interval of $\mathbb{CL}(L)$ is upper semimodular, an upper bounded homomorphic image of a free lattice, and meet semidistributive.*

# References

[1] Dunn, J., Gehrke, M., Palmigiano, A.: Canonical extensions and relational completeness of some substructural logics. J. Symb. Logic **70**, 713–740 (2005)

[2] Erné, M.: Adjunctions and standard constructions for partially ordered sets. In: Hölder, Pichler, Tempsky (eds.) Contributions to General Algebra, vol. 2, pp. 77–106 (1983)

[3] Freese, R., Ježek, J., Nation, J.B.: Free Lattices, *Mathematical surveys and Monographs*, vol. 42. Am. Math. Soc., Providence (1995)

[4] Gehrke, M., Jansana, R., Palmigiano, A.: $\Delta_1$-completions of a poset. Order (to appear). Preprint available from http://www.math.ru.nl/∼mgehrke/GeJaPa.pdf

[5] Haskins, L., Gudder, S.: Semimodular posets and the jordan-dedekind chain condition. Proc. Am. Math. Soc. **28**, 395–396 (1971)

[6] Jónsson, B., Tarski, A.: Boolean algebras with operators I. Am. J. Math. **73**, 891–939 (1951)

[7] MacNeille, H.: Partially ordered sets. Trans. Am. Math. Soc. **42**, 416–460 (1937)

[8] Nation, J.B., Pogel, A.: The lattice of completions of an ordered set. Order **14**, 1–7 (1997)

[9] Ward, M.: The closure operators of a lattice. Ann. Math. **43**, 191–196 (1942)